

REFERENCES

1. Dimopoulos M, Bringhen S, Anttila P, et al. Isatuximab as monotherapy and combined with dexamethasone in patients with relapsed/refractory multiple myeloma. *Blood*. 2021;137(9):1154-1165.
2. Alexanian R, Dimopoulos M. The treatment of multiple myeloma. *N Engl J Med*. 1994;330(7):484-489.
3. Alexanian R, Dimopoulos MA, Delasalle K, Barlogie B. Primary dexamethasone treatment of multiple myeloma. *Blood*. 1992;80(4):887-890.
4. Martin T, Strickland S, Glenn M, et al. Phase I trial of isatuximab monotherapy in the treatment of refractory multiple myeloma. *Blood Cancer J*. 2019;9(41):1-10.
5. Boyle EM, Leleu X, Petillon MO, et al; IFM2014-04 investigators. Daratumumab and dexamethasone is safe and effective for triple refractory myeloma patients: final results of the IFM 2014-04 (Etoile du Nord) trial. *Br J Haematol*. 2019;187(3):319-327.

DOI 10.1182/blood.202009268

© 2021 by The American Society of Hematology

IMMUNOBIOLOGY AND IMMUNOTHERAPY

Comment on Pignarre et al, page 1166

The origin of preplasmablastic cells

Jerome Moreaux | CHU Montpellier; Institute of Human Genetics; University of Montpellier UFR Medicine; Institut Universitaire de France


In this issue of *Blood*, Pignarre et al characterize the genomic events involved in the cell fate decision between activated B cells and plasmablasts.¹ Plasma cells (PCs) play an important role in humoral immunity by synthesizing and secreting antibodies.² Understanding the biological processes that control the production of PCs is critical to both ensure efficient immune response without autoimmunity or immune deficiency and prevent tumorigenesis. The production of interleukin-4 (IL-4) by follicular helper T cells drives B-cell amplification and maturation.³ However, the full molecular mechanisms behind these functions are not fully understood.

Pignarre et al report new biological events driving normal B- to plasma-cell differentiation. Using an in vitro model, naive

B cells were cultured in a 2-step process, which results in differentiation into plasmablasts,⁴ and the authors demonstrated

that cells are destined to differentiate into PCs if there is an early response to IL-4, which results in downregulation of the CD23 cell-surface protein and IL-4/STAT6 signaling. However, B cells maintaining IL-4 signaling did not differentiate. Furthermore, the differentiation of CD23⁻ cells is associated with CBLB E3 ubiquitin ligase downregulation, coinciding with IRF4 induction and with specific chromatin and transcriptional modifications (see figure). The changes were identified by ATAC sequencing and hydroxymethylation profiling. However, no major changes in expression of epigenetic factors were noted. CBLB is known to prevent premature germinal center (GC) exit promoting IRF4 degradation in light zone B cells.⁵ Pignarre et al reported potential STAT6 binding sites in the CBLB promoter, suggesting potential direct regulation, hence the interest in characterizing STAT6 targets, using chromatin immunoprecipitation. CD23⁻ B cells, postactivation, have the characteristics of preplasmablasts with a significant increase in chromatin accessibility at immunoglobulin heavy chain coding loci. Full transcriptomic characterization of the proposed model at a single-cell level would be particularly useful in deciphering the heterogeneity and transcriptional trajectories during B- to plasma-cell differentiation.

The major transcriptional and epigenetic changes reported by Pignarre et al may be associated with changes in nuclear organization during terminal B-cell differentiation. Gene regulation depends on the 3-dimensional chromatin organization



After activation, B cells that are committed to differentiate into PCs downregulate the CD23 cell-surface protein, IL-4/STAT6 signaling, and CBLB activity concomitantly with IRF4 induction. B cells that maintain the IL-4 signaling will not differentiate.

and its regulatory elements. Recent data obtained in a mouse model revealed that B- to plasma-cell differentiation is associated with major changes in chromosome topology that could be driven by PC biology or reflect enhancer-induced modifications in chromatin organization.⁶ B- to plasma-cell transition is associated with compartmentalization changes along with gain in genomic interactions across the Prdm1 locus, increasing genomic interactions between promoter regions and regulatory elements, concurrent with transcriptional induction. In contrast, the early B-cell factor 1 (Ebf1) locus repositions to pericentromeric heterochromatin in association with transcriptional repression. Furthermore, the interchromosomal hubs reported during B- to plasma-cell maturation are associated with histone marks that define transcriptionally active or repressive hubs. The epigenetic landscape characterization together with nuclear architecture study of the human B- to plasma-cell differentiation model developed by Pignarre et al may provide important findings for the understanding of the molecular mechanisms driving B- to plasma-cell fate.

IL-4 production by T-follicular helper and STAT6 mutations activates and drives the IL-4/STAT6 axis in follicular lymphoma.⁷ Recently, single-cell RNAseq characterization of purified GC B cells (GCBs) generated a new single-cell cell of origin classification that identified distinct prognostic subgroups within the GCB and activated B-cell-like diffuse large B-cell lymphoma subgroups.⁸ The analysis of these single-cell transcriptomic resources derived from GC purified B cells, in light of the new data provided by Pignarre et al, may be of particular interest. The results provided by Pignarre et al provide new insights in the molecular mechanisms driving follicular lymphoma biology.

Conflict-of-interest disclosure: The author declares no competing financial interests. ■

REFERENCES

1. Pignarre A, Chatonnet F, Caron G, Haas M, Desmots F, Fest T. Plasmablasts derive from CD23⁻ activated B cells after the extinction of IL-4/STAT6 signaling and IRF4 induction. *Blood*. 2021;137(9):1166-1180.
2. Shapiro-Shelef M, Calame K. Regulation of plasma-cell development. *Nat Rev Immunol*. 2005;5(3):230-242.

3. Crotty S. T follicular helper cell differentiation, function, and roles in disease. *Immunity*. 2014;41(4):529-542.
4. Le Gallou S, Caron G, Delalay C, Rossille D, Tarte K, Fest T. IL-2 requirement for human plasma cell generation: coupling differentiation and proliferation by enhancing MAPK-ERK signaling. *J Immunol*. 2012;189(1):161-173.
5. Li X, Gadzinsky A, Gong L, et al. Cbl ubiquitin ligases control B cell exit from the germinal-center reaction. *Immunity*. 2018;48(3):530-541.e536.
6. Bortnick A, He Z, Aubrey M, et al. Plasma cell fate is orchestrated by elaborate changes in genome compartmentalization and interchromosomal hubs [published correction].

appears in *Cell Rep*. 2020;31(13):107876]. *Cell Rep*. 2020;31(1):107470.

7. Amé-Thomas P, Hoeller S, Artchounin C, et al. CD10 delineates a subset of human IL-4 producing follicular helper T cells involved in the survival of follicular lymphoma B cells. *Blood*. 2015;125(15):2381-2385.
8. Holmes AB, Corinaldesi C, Shen Q, et al. Single-cell analysis of germinal-center B cells informs on lymphoma cell of origin and outcome. *J Exp Med*. 2020;217(10):e20200483.

DOI 10.1182/blood.202009746

© 2021 by The American Society of Hematology

LYMPHOID NEOPLASIA

Comment on Corre et al, page 1192

Deletion 17p: a matter of size and number?

Sonja Zweegman and Niels W. C. J. van de Donk | Amsterdam UMC

In this issue of *Blood*, Corre and colleagues provide solid evidence for a profound negative impact of del(17p) in patients with myeloma who have del(17p) in >55% of their plasma cells. This increased risk is independent of the presence of a TP53 mutation.¹

It is generally accepted that loss of the short arm of chromosome 17 [del(17p)], as determined by fluorescence in situ hybridization (FISH) analysis, is the most important high-risk factor in multiple myeloma, negatively impacting both progression-free survival (PFS) and overall survival (OS).^{2,3} The loss of the TP53 gene, encoding the tumor suppressor protein p53, is supposed to underlie this dismal outcome.⁴ There are limited and conflicting data on the impact of mutations in the TP53 gene, either to a single allele or when associated with del(17p), so-called double hit disease. In a recent DNA sequencing analysis of ~800 heterogeneously treated patients, including both transplant-eligible and noneligible patients, performed by Walker and colleagues, a dismal outcome was found only in patients with biallelic TP53 inactivation with a median OS of 20.7 months. In contrast, copy number loss of 17p only or 1 mutation in TP53 only lacked prognostic impact, showing a similar outcome as compared with patients with TP53 wild type.⁵

Corre and colleagues used a different approach in this study. They first identified 121 newly diagnosed multiple myeloma patients (NDMM) with a del(17p) in >55% of plasma cells who were uniformly treated with intensive therapy, including an autologous stem cell transplantation (ASCT). One-third of these patients had an additional mutation in TP53. As in the study by Walker and colleagues, median OS in biallelic disease was short, only 36 months. However, OS was also significantly worse in patients with del(17p) without TP53 mutation(s), compared with patients lacking del(17p) (52.8 vs 152.2 months, respectively). Therefore, the study of Corre and colleagues supports the continued use of FISH analysis to identify high-risk patients with a poor prognosis based on the presence of del(17p) in >55% of myeloma cells, without the need for additional genome sequencing. This is important, as FISH analysis is widely available and standardized. In addition, FISH data are available from contemporary clinical studies, whereas DNA sequencing data, and thus TP53 status, are generally missing.