


INTRODUCTION & AIM

Multiple myeloma (MM) is the second most common hematological malignancy. Over the past decade, the development of new therapeutics, particularly the immune-based therapies, significantly improved the life quality and survival of patients.

However, enhancing our comprehension and developing novel approaches to reduce the drug resistance associated with anti-CD38 immunotherapies is crucial for patient care. Among factors that could be involved in MM cell resistance, we focused on mitochondrial metabolism, already described as a significant factor influencing treatment response across various cancers.

METHOD

Detection of the mitochondrial respiration by measurement of the Oxygen consumption rate (OCR) and the glycolytic activity by measurement of the Extracellular acidification rate (ECAR) with the Seahorse XF96 Mito stress kit using the optimal cell seeding density



Generation of the Gene expression profile (GEP)-based metabolic index based on 112 genes included 29 glycolytic genes and 83 OXPHOS genes.

$$\text{Metabolic Index} = \sum(\text{Glycolysis genes standardized expression}) - \sum(\text{OXPHOS genes standardized expression})$$

CONCLUSION

Taken together, our findings demonstrate that dysregulation of metabolism holds prognostic significance in newly diagnosed MM patients. Besides, we also reported a link between MM cell metabolism, CD38 expression, and response to anti-CD38 MoAb treatment.

CONTACT INFORMATION

Alizée Steer, Project Manager
a.steer@diag2tec.com

Angélique Bruyer, CEO
a.bruyer@diag2tec.com

Scan to connect



RESULTS

1. GEP-based metabolic index is representative of the heterogeneous functional metabolic activities in Human multiple Myeloma cell lines (HMCL).




Figure 1: Heterogeneity of the metabolic profiles in HMCLs. A. Basal, ATP-linked, Maximal and Spare OCR are heterogeneous in the 25 HMCLs. B. Basal ECAR as well as glycolytic capacity are heterogeneous in the 25 HMCLs. C. Calculated glycolytic and mitochondrial ATP-production of the 25 HMCLs. D. Generation of Metabolic index in 40 HMCLs based on 112 genes including

29 glycolytic genes and 83 OXPHOS genes. E. Correlation between functional Maximal mitochondrial Respiration of 25 HMCLs with GEP-Based metabolic index. F. OCR representation of the 5 HMCLs with the higher metabolic index compared to the 5 HMCLs with the lower metabolic index. G. HMCLs were classified by the quartile method into low, intermediate and high index. HMCLs with a higher metabolic index have higher maximal mitochondrial respiration than HMCLs with a lower metabolic index. * $p < 0.05$ and **** $p < 0.001$ analyzed by one-way ANOVA test followed by Tukey's test (means \pm SD).

2. High metabolic score values are associated with poor outcomes in MM patients.





Figure 2: High metabolic index values are associated with a poor outcome in MM patients. A and C. Representation of the Metabolic index in the patient from the CoMMPass (A) and Montpellier (D) Cohorts. B and D. Patients of the CoMMPass (n=674) (B) and the Montpellier (n=102) (D) cohorts were ranked according to their increased metabolic index. B. The maximum difference in overall survival (OS) was obtained using MaxStat with a cut off 0.07, splitting patients into high-risk (n = 217; red curve) and low-risk (n = 457; green curve) groups. D. The maximum difference in OS was obtained using MaxStat with a cut off -0.11, splitting patients into high-risk (n = 65; red curve) and low-risk (n = 37; green curve) groups.

3. MM patients with low metabolic score show better anti-CD38 Daratumumab treatment responses after relapse and better survival.




Figure 3: GEP-Based metabolic index as potential prognostic value for Daratumumab treatment. A. MM patients were considered "responders to Dara" if they obtained at least a partial response as a best response to Dara. MM patients were considered "non-responders to Dara" if they had stable or progressive disease. ** $p < 0.01$ was analyzed by two-tailed t-test (means \pm SD). B. Patients of the Montpellier cohort treated with Dara after relapse (n=42) were ranked according to their increased metabolic index. C. The maximum difference in overall survival (OS) was obtained using MaxStat with a cut off -0.05, splitting patients into high-risk (n = 17; red curve) and low-risk (n = 10; green curve) groups.

4. Low CD38 cell surface expression correlates with high functional metabolism in HMCLs and high metabolic score in MM patients.




Figure 4: CD38 expression correlates with functional metabolism and metabolic score in HMCLs and MM patients. A. Correlation between functional basal mitochondrial Respiration of 23 HMCLs with their respective CD38 expression measure using surface antigen binding capacity by flow cytometry. B. CD38 expressions were measured on 24 MM patients by flow cytometry (median fluorescence intensity/MFI). Patients were separated into 2 groups (low and high expression) via their CD38 expression MFI mean. *** p value < 0.001 and **** p value < 0.0001 were analyzed by two-tailed t-test (means \pm SD).